

WHAT'S THE "WILD" SHRUB WITH BRIGHT RED BERRIES?

People from this part of the state and on east have been reporting shrubs with bright red berries growing wild. The berries are clustered around the stem and the leaves are still a bright green color. These are likely one of two species of bush honeysuckle, (Amur or Tartarian), which can get 6-20 feet tall. This landscape shrub has become a serious understory invasive throughout the Midwest from eastern Kansas to Ohio. Many states have it on their noxious weeds list. All of our native honeysuckles are vines, similar to the vining Japanese honeysuckle. Bush honeysuckles are also noticeable in the spring as they put out leaves much earlier than most other trees and shrubs. Leaves also stay green much later into the fall. This long growing season gives it a competitive advantage over other native species, and the vigorous growth can take over a woodland understory, reducing the number of native woodland wildflowers and other shrubs. Hunters and hikers are all too familiar with these path choking, understory dominating masses of foliage. If you want to promote native species on your property, and also not get your area so choked up with growth that you can't walk through it, then controlling bush honeysuckles is a must. Honeysuckle seedlings can be readily hand pulled when the soil is damp. Chemical control is needed for larger infestations, as cutting alone results in vigorous resprouting. Foliar applications of glyphosate (i.e., Roundup) in late summer and fall works well as does applications of Crossbow (2,4-D + triclopyr). Treating cut stumps with Tordon RTU (picloram), or concentrated (20% - 50%) glyphosate is also quite effective. Several studies have shown basal spraying with triclopyr (Garlon) not to be effective, while basal applications with 2,4-D or picloram products work well, using an oil carrier to penetrate the bark. Cut stump and basal treatments can be done when the areas to be sprayed are dry and not frozen. Please follow all label instructions when using pesticides.



Horseradish

Horseradish is ready to dig after a hard freeze kills the foliage (usually November or December). The large roots can be harvested while smaller, pencil sized roots can be cut in 6-8-inch-long sections as 'seed' or 'sets' for next year's crop which are then immediately re-planted. Another option is to leave the horseradish in the ground and dig as needed. If you choose the latter option, be sure to heavily mulch the area so that the ground doesn't freeze. To use horseradish, peel the large, fleshy roots and cut into sections. Use a blender or food processor to chop the roots along with a small amount of water and a couple of ice cubes. Vinegar or lemon juice is added to stop the process that produces the "bite" of horseradish. Add immediately after blending for a mild flavor or wait up to 3 minutes to give the horseradish more kick. Use 2 to 3 tablespoons of vinegar or lemon juice per cup of horseradish sauce along with ½ teaspoon of salt for flavor. Horseradish has an extremely strong odor and so you may wish to open the blender or food processor outdoors and to keep your face away from the container when opening. Store ground horseradish in a tightly sealed jar in a refrigerator until ready for use.

Ice Melters

There are five main materials that are used as chemical de-icers: calcium chloride, sodium chloride (table salt), potassium chloride, urea, and calcium magnesium acetate. Calcium chloride is the traditional ice-melting product. Though it will melt ice to about -25 degrees F, it will form slippery, slimy surfaces on concrete and other hard surfaces. Plants are not likely to be harmed unless excessive amounts are used. Rock salt is sodium chloride and is the least expensive material available. It is effective to approximately 12 degrees F but can damage soils, plants and metals. Potassium chloride can also cause serious plant injury when washed or splashed on foliage. Both calcium chloride and potassium chloride can damage roots of plants. Urea (carbonyl diamide) is a fertilizer that is sometimes used to melt ice. Though it is only about 10% as corrosive as sodium chloride, it can contaminate ground and surface water with nitrates. Urea is effective to about 21 degrees F. Calcium magnesium acetate (CMA), a newer product, is made from dolomitic limestone and acetic acid (the principal compound of vinegar). CMA works differently than the other materials in that it does not form a brine like salt but rather helps prevent snow particles from sticking to each other or the road surface. It has little effect on plant growth or concrete surfaces. Performance decreases below 20 degrees F.



Limited use of any of these products should cause little injury. Problems accumulate when they are used excessively and there is not adequate rainfall to wash or leach the material from the area. Since limited use is recommended it is best to remove the ice and snow by hand when possible. When they are applied, practice moderation. Resist the temptation to over apply just to make sure the ice and snow melts. Keep in mind this can damage concrete surfaces as well as the plants and grass growing along the walks and driveways. These problems are normally latent and do not show up until spring or summer.